NOTES AND REVIEWS

Sverre Petterssen. Weather Analysis and Forecasting. New York (McGraw-Hill Book Co.), 1940. 505

pp., 249 figs.

The purpose of this book is to assemble the practical methods now available (most of them developed comparatively recently) for forecasting weather on the basis of physical principles. A general knowledge of meteorology is presupposed, and only those aspects of meteorology are treated which experience has shown to be directly applicable to actual forecasting. The book embodies the original theoretical investigations and the accumulated experience of the author in synoptic meteorology over a long period of years. The author's methods are distinguished by the introduction of precise formulations of synoptic concepts, in terms of quantities that can be measured in ordinary meteorological observation.

The first chapter contains a brief treatment of 17 different quantities in terms of which the physical characteristics of air masses may be described. It is followed by a chapter on the theory of atmospheric stability, convec-

tive phenomena, and fog, and the uses of various thermodynamic diagrams. Chapter III discusses the source regions, production, classification, and transformation of air masses. Two chapters are then devoted to the fundamental dynamical equations of motion, the gradient wind, kinematic analysis, frontogenesis, and the characteristic phenomena of fronts, followed by a chapter on waves and the wave theory of cyclones. Chapter VIII is on isentropic analysis, and was written by Jerome Namias. chapters on the kinematic forecasting of the displacements of pressure systems, fronts, and air masses, and the forecasting of deepening and filling, are followed by the concluding chapter on the actual technique of analysis and forecasting, with several examples.

EDMUND SCHULMAN. A Bibliography of Tree-Ring Analysis. *Tree-Ring Bulletin* (University of Arizona, Tucson), Vol. 6, No. 4, April 1940. This entire issue of the *Tree-Ring Bulletin* is devoted to a 12-page bibliography comprising 412 titles on tree-ring analysis.

METEOROLOGICAL AND CLIMATOLOGICAL DATA FOR AUGUST 1940

[Climate and Crop Weather Division, J. B. KINCER in charge]

AEROLOGICAL OBSERVATIONS

By EARL C. THOM

The mean surface temperatures during August (Chart I) were above normal over most of the United States; temperatures were slightly below normal, however, over considerable of the western part of the Mississippi River Valley area, and a small area in northeastern Texas had temperatures from 2° to 3° (F.) below normal. Temperatures were also below normal over the northern Atlantic States, with temperatures about 3° below normal in New Jersey and eastern New York. The warmest area for the month, relative to the normal, was Montana, western Wyoming, northern Utah, and northern Nevada with mean temperatures about 5° above normal in most of this area.

At the 1,500-meter level the direction of the resultant winds for the month (chart VIII) was considerably to the north of normal (clockwise turning) over the Atlantic Coast States and also over eastern Texas. The direction of the resultants for the month had the opposite turning over the Great Lakes and the east central states. The same general diversion from normal in the direction of the resultant winds for August occurred at the 3,000-meter level as at the 1,500-meter level, except that the shift of direction to northward was more pronounced over eastern Texas and extended over Oklahoma. Based on data for the small number of stations for which 5 a.m. normal records are at hand, the direction of the resultant wind for the month at 5 p. m. was south of the corresponding 5 a. m. normal over the eastern half of the country and over the extreme northwest with the opposite shift occurring over the remainder of western United States.

The 5 a. m. resultant velocity for August was in general below the normal resultant velocity over the United States at both the 1,500-meter and the 3,000-meter levels. resultant velocity was about 3 meters per second below normal at the 1,500-meter level along the northern Atlantic coast and at Fargo, N. Dak., while the corresponding velocity was slightly over 4 meters per second below normal for the 3,000-meter level at Fargo and Sault Ste. Marie. At the 5,000-meter level the 5 p. m. resultant normal was considerably above the corresponding 5 a.m. velocity over the South Central States and was considerably below this normal over the Northeast and North

Again in August the areas having a decided shift in direction of the resultant winds from normal, at both the 1,500-meter and the 3,000-meter levels, agree well with the areas having corresponding surface temperature departures for the month.

At the 1,500-meter level the direction of the 5 p. m. resultant for the month was south of the corresponding 5 a. m. resultant (counterclockwise turning) over a large part of the United States. The opposite shift in resultant wind direction at this level occurred over the Great Lakes and part of the North Atlantic States, over part of the Southeastern States and over most of the extreme western and extreme southwestern portions of the country. At the 3,000-meter level the shift of the resultant wind to the northward from 5 a.m. to 5 p.m. was noted over larger areas, this shift having occurred over the Gulf coast, southern Atlantic coast, part of the east Central States, over the Pacific coast and most of the Southwest and West Central States.

The changes in resultant velocity at 5 p. m. from those at 5 a. m. were well distributed at both the 1,500-meter and the 3,000-meter levels. At the 1,500-meter level, however, the West Central, the Central, and the Southeastern States all showed lower resultant velocities at 5 p. m. than at 5 a. m., the velocity departures in this area amounting to a decrease of Beaufort force one or

It is of interest to note that at Las Vegas, Nev., the pilot balloon observations attained a height of 18 kilometers or over on 16 days during the month and that the resultant wind, based on those observations was 118° (ESE), 2.0 meters per second for that level.

The upper air wind data discussed above are shown in

table II and in charts VIII to XI.

At all levels from 3,000 meters up to at least 17,000 meters over the United States (table I) the highest mean monthly pressures for the month of August occurred over the extreme southwestern part of the country; the maximum occurred at Phoenix at most of these levels. At

several of the levels above 8,000 meters, however, the mean pressure at other stations (Brownsville, San Diego, or El Paso) were the same as at Phoenix. The lowest mean monthly pressure occurred at Sault Ste. Marie at the 15,000-meter level as well as at all levels from 3,000 meters to 13,000 meters inclusive. At the 14,000-meter level the same minimum pressure, 151 mb., was recorded at Great Falls, Mont., and at Sault Ste. Marie, Mich., while 110 mb. pressure was recorded at three stations, Great Falls, Joliet, and Sault Ste. Marie, at the 16,000-meter level.

Mean monthly pressures were the same or lower in August than in July at most stations for all levels below 5,000 meters. At higher levels mean pressures were higher in August than in July over the eastern half of the United States and along the Pacific coast with little change over the rest of the United States. The station showing the largest increase in mean pressure at upper levels from July to August was Sault Ste. Marie, this increase being 4 mb. at 8,000 meters, 6 mb. at 11,000 meters, and 5 mb. at 14,000 meters over this station.

The greatest difference between the mean maximum pressure for August and the corresponding mean minimum was 12 mb. which occurred at the 8,000-meter level. The steepest pressure gradient at this level occurred between Lakehurst and Washington where a difference of 3 mb. was noted along a horizontal distance of less than 200 miles. Another steep pressure gradient, a difference of 5 mb., occurred between Sault Ste. Marie and Joliet.

At levels below 8,000 meters the mean temperatures were in general higher in August than in July over the Great Lakes and East Central States and along the Pacific Coast and were lower than in July over the balance of the country. Almost without exception mean temperatures were higher than in July at all thousand-meter levels from 9,000 meters to and including 13,000 meters. At the 14,000-meter level four stations in the eastern half of the country and one in the western half had mean temperatures in August lower than those in July. At levels from 15,000 meters to 18,000 meters inclusive, mean temperatures were generally lower than in July except that in the southwest temperatures were higher. At El Paso, for example, the temperature at these four levels averaged nearly 2° C. higher than in the previous month.

At levels below 5,000 meters mean temperatures were generally lower than in August of 1939 except over the Southeast, the Gulf Coast and over the West Central States where small increases in temperature from those of the preceding year were noted. At most of the thousand-meter levels from 5,000 to 14,000 meters inclusive, temperatures were lower than in 1939 at Oakland and at Joliet and were higher than 1939 at Denver and El Paso. At most other stations no marked temperature changes from the previous year were evident.

The maximum altitude of the level of mean freezing temperature for August was 5,100 meters over Phoenix and the minimum altitude of this level was 3,700 meters over Sault Ste. Marie. The altitude of mean 0° C.

temperature was 4,900 meters or higher at all Weather Bureau Stations south of 35° N. latitude, while the corresponding altitude ranged between 4,400 meters and 4,800 meters for such stations in the United States between 35° and 45° latitude and was the same or lower for all stations north of this latitude belt. The altitude of the level of average freezing temperature was several hundred meters higher than in July over the eastern half of the country, the maximum increase of 800 meters occurring over Sault Ste. Marie.

The lowest free-air temperature at standard one thousand meter levels in August was -82.6°C . $(-116.7^{\circ}\text{ F.})$ recorded at 16,000 meters over Brownsville, Texas. Minimum temperatures of -70° C. (-94° F.) or lower were recorded at all radiosonde stations in the United States south of 35° N. latitude while the minimum recorded at all stations (except Lakehurst) north of this line was -70° C. or higher, with the highest minimum observed in the United States, -63.2° C. $(-81.8^{\circ}\text{ F.})$

occurring at Bismarck, N. Dak.

Table 3 shows the maximum free-air velocities and their directions for various sections of the United States during August, as determined by pilot-balloon observations. The extreme maximum for the month was 53.6 meters per second (119.9 miles per hour) observed over Huron, S. Dak. on August 5, 1940. This high wind was blowing from the west at an elevation of 9,540 meters (nearly 6 miles) above sea level. A maximum, 58 meters per second, for the month of August occurred last year over Huron, S. Dak. at 13,740 meters above sea level. In August 1938 a still higher maximum wind, 69.8 meters per second, was observed at an elevation of 2,660 meters over Havre, Mont.

Tropopause data for August showing the mean altitude and temperature of the tropopause at various stations are shown in table 4 and on chart XIII.

MEAN ISENTROPIC CHART 1

Two anticyclonic eddies are the most prominent features of the mean isentropic chart $(\theta=314^{\circ})$ for August 1940 (chart XII). The eastern cell is displaced southward of its usual position, resulting in above normal precipitation in the Middle and South Atlantic States, but with large deficiencies in New York and New England. Downslope motion is indicated by the resultant winds in the region west of the Rockies, where little or no precipitation occurred; but over the upper Mississippi Valley marked upslope components are shown, and as a consequence the precipitation was above normal. The large excesses in precipitation in Louisiana, as well as those in North Carolina and Virginia, were due largely to tropical cyclones the effect of which was not prolonged enough to show up on the mean moisture pattern.

The data for Medford, Oreg. were neglected in constructing chart XII because the mean there was based on only 18 observations.

¹ Prepared by the Research Division.

Table 1.—Mean free-air barometric pressure in millibars, temperature in degrees Centigrade, and relative humidities in percent, obtained by airplanes and radiosondes during August 1940

									puan																						
										8	tations	and	eleva	tion	as in n	ieters	abo	ve se	a le	vel											
Altitude (meters)	Bism		t, N. D 5 m.)	ak.	Bro	wnsv (6 m.	ille, T	еж.	Ch		on, S. m.)	c.	C	oco	Solo, (15 n		,1 1			nver, 1,616	Colo. m.)				o, Te 3 m.)				ly, N ,908 1		
m. s. l.	Num- ber of obs.	P.	т.	R. H.	Num- ber of obs.	P.	т.	R. H.	Num- ber of obs.	Р.	т.	R	Nur ber obs	of	Р.	т.	R. H.	Nu ber ob		Р.	т.	[뜹· b	Num- er of obs.	P.	т.	R	Nur ber obs	of 1	Р.	т.	R. H.
Surface	31		18. 2		27 27	1, 013 958	24.	5 87	30	1, 01 96	0 22.	9 7	79	17	1, 013 958	25. 24.	2 9	4	31	842	18. 3	62	21	882	23.	6 4	19	31 8	311	17. 5	28
1,000 1,500 2,500 3,000 4,000 5,000 6,000 7,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 12,000 11,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000	31 31 31 31 31 31 31 31 30 28 28 22 24 24 24 21 13 5 7	111 94 80 69 58	18. 0 15. 1 12. 2 9. 0 2. 9 -3. 8 -17. 4 -17. 4 -25. 1 -32. 8 -47. 3 -47. 3 -52. 7 -59. 1 -59. 1 -59. 1 -59. 1 -59. 1	53 53 53 46 40 36 32 31 29	27 27 27 27	158 134 113 95 81 68	20. 17. 15. 6. -11. -18. -26. -33. -41. -48. -56. -62. -68. -71. -67. -64. -61.	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	30 30 30 30 30 30 30 29 5 5 28 28 28	15 13 11 9 8 6 5	5 17. 15. 16. 16. 17. 16. 17. 17. 17. 17. 17. 17. 17. 17. 17. 17	3 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	74 74 76 70 88 86 66 66 67 77 67 66 66 66 66 66 67 77 77	17 17 16 15 15 11 11	904 854 805 759 714 633	22. 19. 17. 14. 11. 6.	1 9:6 9 1 8:3 3 8:8 8 8:0 7:	1 6 4 2	31 31 31 31 29 28 28 28 29 28 29 28 20 25 22 19 18 6	433 - 379 - 830 - 286 - 212 - 182 - 155 - 112 - 95 - 81 - 69 -	19. 2 16. 2 12. 9 5. 7 -9. 0 -15. 2 -22. 0 -51. 5 -61. 5 -64. 3 -65. 1 -65. 1 -65. 1 -65. 1 -57. 4 -55. 6		211 211 220 220 220 220 220 220 220 220	332 288 249 215 184 157 133 113 95 81	17. 13. 6. -12. -19. -26. -34. -42. -50. -62. -67. -69.	8 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	944 88 33 36 66 33 33 32 11 90	31 7 31 7 31 6 31 6 31 4 31 4 31 31 3 31 331 3 31 331 3 31 329 229 1 229 1 229 1 229 1 229 1	194 133 — 133 — 286 — 246 — 212 — 181 — 154 — 112 — 95 — 80 — 69 —	20. 2 20. 1 16. 4 7. 6 -9. 7 -15. 8 -22. 9 -38. 5 -45. 7 -61. 8 -61. 2 -64. 6 -64. 8 -61. 0 -65. 7 -65. 4	25 24 24 28 38 45 42 38 35 32
Altitude	Grea	t Fa (1,1	lls, Mo 17 m.)	ont.	;	Joliet, (178		T	Lake		,N.J.			edfo	ord, On		Τ	Nasl		 е, Те	 nn,	7	Norfol	lk, V 0 m.)			0		nd, (2 m.)	Calif.	
(meters) m. s. l.	Num- ber of obs.	Ρ.	т.	R. H.	Num- ber of obs.	P.	т.	음: [1	Num- per of obs.	Р.	т.	R. H.	Num- ber of obs.	P	. Т.	R	· be	nm- er of bs.	P.	т.	R. H.	Num ber o obs.	P.		т.	R. H.	Num- ber of obs.	P		т.	R. H.
Surface 500 1,000 1,500 2,000 2,500 3,000 4,000 5,000 6,000 10,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 12,000 11,000 11,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000	31	490 429 375 325 282 242 207 177 151 129 110 94 80 68	2 20.4 3 17.5 7 13.6 9 4 2 2.2 2 2.2 7 -5.0 0 -11.0 0 -11.0 0 -15.6 6 3-43.1 2 -50.5 6 3-43.1 -59.0 0 -58.8 0 -58.8 0 -58.8 0 -58.8 0 -58.8 0 -58.8 0 -58.8	322 33 35 35 38 44 42 40 40 38	29 29 29 28 28 28 27 26 26 26 26	209 - 178 - 152 - 130 - 110 - 93 - 79 - 68 - 57 -	19. 3 20. 7 18. 7 15. 8 10. 1 7. 0 1. 4 4. 4 3 10. 3 11. 3 10. 3 11. 3 12. 4 4. 4 3 31. 9 9 446. 9 52. 5 6 6 6 6 7 7 7 7 7 7 8 9 8 9 9 1 9 1 9 1 9 1 9 1 9 1 9 1 9 1		31 31 31 31 31 30 30 30 30 30 29 29 29 29 29 29 29 29 29 29 29 29 29	283 245 210 179 153 130 111 95 81 68	17. 0 8 14. 8 12. 4 10. 3 6. 5 6. 16 -3. 7 -9. 7 4 -23. 5 -31. 0 -35. 2 -45. 8 -62. 9 -60. 8 -58. 2 -62. 9 -64. 5 -58. 4 -64. 5 -58. 4 -64. 5 -58. 4		19 19 19 19 19 19 19 19 19 18 17 17 17 17 17 15 15 15 15 15 14 14 14 12 18 18 18 18 18 18 18 18 18 18 18 18 18	955 90 855 80 77 63 555 48 43 37 32 24 21 11 11 11 11	566 22 202 20 511 17 512 15 566 14 122 11 388 -22 381 -15 381 -15 384 -40 455 -47 380 -56 53 -59 380 -61 111 -63 380 -61	.79 4 4 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	 	31 31 31 31 31 31 31 31 31 29 27 26 26 26 25 25 21 17 14 9	995 959 906 855 806 759 715 634 434 434 380 228 218 115 80 68 58 50	22. 23. 21. 17. 14. 11. 9. 41612193442566367645956555656.	5	2 2 2 2 2 2 2 2 2 2	7 97 87 87 87 77 77 66 5 5	63 09 557 508 661 116 334 559	22. 8 21. 4 18. 5 15. 4 12. 1 9. 4 12. 1 8. 3. 8	87 79 74 72 69 63 56 47 39	31 31 31 31 31 31 31 31 31 31 31 31 31 3	888888888888888888888888888888888888888	957 902 903 903 950 950 950 950 950 950 950 950	15. 6 7 7 121. 1 1 18. 0 1 11. 8	

See footnotes at end of table.

Table 1.—Mean free-air barometric pressure in millibars, temperature in degrees Centigrade, and relative humidities in percent, obtained by airplanes and radiosondes during August 1940—Continued

							-			s	tation	s and	l eleva	tions i	mete	rs abo	ove sea	leve	el .										_
Altitude (meters)	Okl	lah cla.	ma Ci (391 m	y,	Or	naha, (301	Nebi m.)		Pearl	Harbo (6 n		H.13	Pe	nsacol (24 i	, Fla. ¹ n.)		Pb	oeni (339	x, Ariz. m.)	,	Sa	n Dieg (19)		dif.¹	8	ault Mich	Ste. 1	Marie 21 m.)	·,
m. s. l.	Num- ber of obs.	Р	. т.	R. H.	Num- ber of obs.	Р.	т.	R. H.	Num- ber of obs.	P.	т.	R. H.	Num- ber of obs.	P.	т.	R. H.	Num- ber of obs.	Р.	т.	R. H.	Num- ber of obs.	P.	T.	.]	R. Nu ber ob	of I	P.	т.	R. H.
Surface 500 1,000 1,500 2,000 2,000 2,500 3,000 4,000 5,000 6,000 7,000 8,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 11,000 12,000 11,000 12,000 11,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000 12,000	31 31 31 31 31 31 31 31 30 30 29 29 29 29 29 27 27 25 24 24 21 18 14	78 71 63 56 49 43 37 33 28 24 21 18 11 18 11 18 11 18 11 18 18 18 18 18	8 23.4 4 22.4 4 19.5 5 16. 9 13.5 5 10. 14 4. 3 -7. 3 -13. 9 -20. 9 -20. 17 -35. 8 -43. 3 -51. 12 -57. 5 -63. 2 -67.	5 70 63 64 64 64 64 64 64 64 64 64 64 64 64 64	31 31 31 31 31 31 30 29 29	490 430 376 327 284 245 211 181 154 131 112 95 81	20. 5 20. 6 19. 2 16. 8 13. 9 10. 9 8. 3 2. 1 -9. 5 -15. 7 -22. 7 -38. 1 -45. 0 -51. 1 -51. 1 -62. 3 -62. 3 -62. 5 -63. 6 -63. 6	75 69 66 64 62 61 61 55 48 44 43 41 41		1,014 959 905 854 805 759 715 633	24. 6 22. 0 19. 1 16. 4 14. 9 13. 1 10. 8 5. 1	84 90 86 71 52 41	31 31 31 31 31 31 31 31 22 7 7 7 6 6 6 6	760 716 634 561 494 484 381 332 288 249	24. 5 24. 1 21. 0 18. 0 15. 2 9. 7 -2. 6 -8. 4 -14. 4 -21. 0 -28. 7 -36. 6 -44. 6	77 79 78 74 65 58 53 56 52 50	311 311 311 311 311 311 311 312 29 29 29 29 29 211 65	953 902 852 805 760 717 636 436 436 436 382 233 290 251 185 185 184 113 96 81 169 588	27. 3 23. 1 18. 8 15. 0 7. 7 0. 8 -5. 6 -12. 0 -18. 9 -26. 6 -34. 3 -42. 1 -49. 8 -56. 9 -62. 8 -67. 2 -68. 8 -68. 7 -66. 3	40 36 34 36 40 47 52 61 63 62 54 47 44 42	20 20 20 20 20 15 12 12 12 12 12	957 903 853 804 758 716 635 561 495 380 331 288 249 213 185 157	18 23 23 21 18 14 6 6 -0 -7 -13 -20 -28 -35 -43 -50 -68 -69	.0 .3 .8 .4 .0 .3 .7 .7 .0 .5 .7 .7 .5 .6 0 .5 .6 0 .5 .6 .6 .6 .6 .6 .6 .6 .6 .6 .6 .6 .6 .6	83 74 36 27 28 228 226 29 227 20 15	31 9 31 8 31 8 31 8 31 7 30 6 30 5 29 4 28 3 27 2 27 2 27 2 27 2 27 2 26 1 25 1 25 1 16	60 06 05 53 63 63 63 63 64 65 66 67 67 67 67 67 67 67 67 67	14. 9 16. 1 14. 5 11. 5 8. 8 6. 2 3. 5 -7. 1 13. 5 220. 2 27. 5 34. 6 41. 6 -52. 0 -55. 4 -55. 4 -55. 8 55. 8 -55. 8	
		<u>-</u>	<u> </u>	<u> </u>	<u> </u>							<u> </u>	<u></u>	<u> </u>							<u> </u>		<u> </u>				<u> </u>		
<u>"</u>				<u>!</u>	!			1	<u> </u>	Stati	ons ar	id ele	vation	s in me	ters at	ove s	sea levo	el					<u> </u>		Lat	e rep	ort Ju	uly 19	40
Altitude (me	eters) p	a. s	. 1.	<u>!</u>	Seattle,	Wasl	1. ¹	St.	Thoma			7	ashin			T	antic S		n No. 1	2 A	l Lianti	e Stati	on No	0. 24		ortlan		/I aine	40
Altitude (me	eters) u	n. 8		Num ber of obs.	(27)	Washm.)	n.¹	St.	n- of P.	m.)	[,1 4	7	ashing (gton, I	o. C. ¹	T	antic S	tatio 5 m.)	n No. 1	N	tlanti	e Statie (5 m.)	on No	R. H.		ortlar (1:	ad, M 9 m.)	Maine	== 40 R. H.

Note.—All observations taken at 12:30 a. m. 75th meridian time, except those at Washington, D. C., and Lakehurst, N. J., where they are taken before 5 a. m., 75th meridian time. At Norfolk, Va., observations are taken at about 6 a. m., and at Pearl Harbor, T. H., after sunrise.

None of the means included in this table are based on less than 15 surface or 5 standard level observations.

Number of observations refers to pressure only, as temperature and humidity data are missing for some observations at certain levels; also, the humidity data are not used in daily observations when the temperature is below -40.0° C.

¹ U. S. Navy. ² Airplane observations. ³ In or near the 5° square:

Lat. 35°00′ N. to 40°00′ N. Long. 55°00′ W. to 60°00′ W.

In or near the 5° square:

Lat. 40°00' N. to 45°00' N. Long. 40°00' W. to 45°00' W.

Airplane observations and radiosondes.

Table 2.—Free-air resultant winds based on pilot balloon observations made near 5 p. m. (75th meridian time) during August 1940. Directions given in degrees from North ($N=380^{\circ}$, $E=90^{\circ}$, $S=180^{\circ}$, $W=270^{\circ}$)—Velocities in meters per second

				y • ·			· · ·				0. 0	\ <u>`</u>						~-		,		~ .								- P									
Altitude		biler Tex 537 n	. '	All N (1,0	ouqu que, . Me 630 1	ex. m.)	i	tlant Ga. 99 n	•	N	illing Aont 095 n	.	N.	mar Da 12 m	k.	3	3oise dah 70 m	0	vil	row le, 1	ex.	1	uffal V. Y 20 m	. 1	to	urlin n, V 32 n	'ŧ.	to	harle 1, 8. 18 m.	C.		hica Ill. 92 n		na	linci ti, O 57 n	hio	- (enve Colo 327 1	
(meters) m. s. l.	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity
Surface	31 31 31 29 28 20 15 14 12 11 10				264 245 272 325 324 319 289 285 297 274 284	0.5 1.3 .6 1.3 2.6 3.8 3.9 5.6 10.0 11.5 8.5 5.4	30 30 29 25 21 20 17 14 11	51 26 38 33 285 288 267 282 283 294 270 289	1. 1 1. 2 1. 1 . 3 . 8 2. 3 3. 0 3. 1 4. 1 5. 7 5. 6			1, 3 1, 8 . 7 1, 2 3, 4 7, 7 9, 7 10, 7 12, 2 15, 2 16, 8	31 28 27 26 25 21 19 18 13 10	70 95 249 271 293 284 299 300 304 298 304	0. 7 . 6 2. 4 2. 8 5. 5 7. 6 8. 0 8. 2 10. 4 12. 3		200	4. 6 3. 7 3. 0 3. 8 5. 0 7. 1 9. 2 9. 4 9. 1 10. 0 14. 5 15. 4	31 30 28 25 24 21 19 14 12	136 138 153 159 169 179 177 210 187 166 351 39	5.9 6.3 4.2 2.7 1.7 1.1 1.7 1.4 .7 .4		79 141 218 235 236 267 280 283 297 290	0. 9 .7 1. 9 2. 4 2. 4 3. 9 3. 8 4. 9 6. 5	31 30 29 28 24 20 18 15 10	207 219 231 252 266 270 264 286 298 298	1. 3 3. 0 2. 3 3. 4 4. 6 4. 8 3. 7 5. 5 7. 2 8. 3	29 29 28 27 25 23 17 12 10	105 112 86 68 289 307 284 276 326	1. 4 2. 8 1. 8 . 7 1. 3 1. 9 2. 2 1. 4 3. 3	31 31 29 28 27 24 22 17 14 11 11	46 86 198 226 234 248 266 264 275 272 264	0. 7 1. 7 2. 3 3. 0 4. 4 4. 9 5. 3 5. 2 7. 0 8. 9 11. 7	31 30 27 24 21 16 10	108 155 180 210 217 237 222 206 271	0. 6 1. 2 1. 6 1. 6 2. 3 2. 6 2. 1 1. 7 1. 7		355 8 9 333 320 320 325 329	1. 5 2. 3 1. 6 1. 4 4. 5 5. 5 7. 1 9. 2 7. 5
	(1	l Pa Tex ,196	so, m.)	El:	y, N 910 1	ev. m.)	Gra tio (1	nd J n, C ,413 1	unc- olo. n.)	Gre	ensb N. C 71 m	oro,]	Iavr Vion 66 m	t.	vi	ickso lle, I 14 m	da.		s Ve Nev 570 i	gas, '. n.)		tle R Ark 79 m	•		edfo Oreg 110 m	.		mi, 10 m		olis	inne , Mi 261 n	nn.	1	Iobi Ala 10 m	·	1 7	shvi Fenn 94 m	١.
Alti.ude (meters) m. s. l.	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity
Surface					216 216 224 222 217 226 249	1. 9 2. 4 2. 8 3. 1 3. 6 4. 8 6. 1 7. 3 14. 5	31 31 31 31 31 28 24 10	352 347 331 286 288 292 295 274		24 24 23 22 19 19 14 12	37 44 94 196 276 297 285 274 286	1.6 1.8 1.1 .3 1.3 3.2 5.0 5.2 6.0	1 1	271 269 216 234 230 237 257 267 269	0. 4 . 9 1. 8 1. 5 4. 6 6. 9 9. 1 12. 3 12. 2	077	114 141 220 269 296 295 267 270 290 311 329 323	2.5 2.5 1.7 1.9 2.3 2.5 2.6 1.9 2.1 2.2 2.1 4.5	31 31 31 31 31 30 28 27 26 23 23 21 20 16	174 184 195 207 205 214 215 221 238 259 258	1		152 146 168 161 144 133 118 298	1.7 1.3 .8 .7 1.1 1.6 4.0	31 31 31 31 31 31 30	246	1.221.234.5679.6577	31 31 31 31 31 31 30 24 22 18 13	112 136 177 226 221 225 216 242 181 117 79	1.8 1.4 .6 .5 1.0 1.3 1.6 1.2 1.6 2.0	26 24 18 17 14 10	92 114 256 239 251 298 272 257	0. 9 1. 0 . 6 1. 4 1. 5 1. 9 3. 5 4. 8	31 31 30 29 27 26 25 20 18 15	170 190 123 9 345 348 292 238 242 243	2. 0 2. 1 .8 .9 .9 .6 .8 2. 9 3. 5 4. 0	31 31 31 31 31 25 21 10	10 28 19 350 303 264 267 291	0.9 .8 .9 .4 .5 1.7 1.9 4.2
	No.	ew Y I. Y. m.)	ork, (15	O Ca	akla lif (8	nd, 3 m.)	Ci	rlaho ty, O 402 n	kla.	1 :	mah Nebr 106 m	•	P)	noen iz. (i m.)	ix, 344	D	Rapi ity, ak. (m.)	d S. 982	St M	. Lo Io. (m.	181	tor	an A io, 7	Геx.	Sar	n Di alif. m.)	ego, (15	1	ult 8 Marid ich. ((m.)	e, 230	8 W	eatti ash. m.)	le, (14	S ₁ W	oka ash. m.)	(603	w tor	ashin ı, D. 10 m	ng- C.
Altitude (meters) m. s. l.	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity	Observations	Direction	Velocity
Surface	29 29 27 25 20 17 13	147 162 225 268 286 299 278		31 30 29 29 29 29 29 29 27 26 22 18 13	281 269 222 200 193 181 186 201 201 210 233 248 238	5. 0 3. 1 3. 3 3. 6 3. 7 3. 6 5. 4 5. 9 7. 8 12. 8 13. 4	31 31 31 32 30 32 32 32 32 32 32 32 32 32 32 32 32 32	158 162 164 167 204 275 292 324 316 298	2. 2 2. 7 2. 4 2. 1 1. 6 1. 9 2. 7 5. 7 6. 7	31 31 30 30 28 27 24 20 18 10	147 145 194 220 247 275 281 290 292 293 287	1. 4 1. 7 1. 8 2. 8 4. 0 5. 6 7. 6 9. 8 10. 1 13. 4	31 31 31 31 31 31 31 31 30 28 26 22 20 13	238 236 247 240 228 213 231 67 89 76 292 272 270 288 157	1. 5 1. 9 2. 3 2. 0 1. 2 . 5 1. 7 1. 7 5. 0 9. 8 5. 3 1. 4	31 31 31 30 29 28 23 22 15 13 12 10	116 137 161 240 288 305 309 303 314 308 294 275	1. 9 1. 9 1. 2 .8 1. 1 4. 5 6. 5 7. 8 10. 0 11. 9 13. 9 15. 9 11. 6 7. 6	30 30 29 28 28 24 22 19 16 14	213 222 201 199 219 265 283 281 286 273	1, 2 1, 1 1, 3 2, 6 3, 0 3, 8 4, 8 5, 9 6, 5 6, 2	31 31 31 31 29 27 25 24 20 20 17	119 116 118 127 135 154 302 299 313 313 327 360	1. 7 2. 0 1. 7 1. 7 .8 .8 .4 1. 5 1. 5 3. 3 1. 5 2. 7	31 30 30 30 30 29 26 24 20 11	266 288 303 296 242 199 180 181 196 219 240	3.9 2.26 2.01.4 1.6 2.3 3.3 3.5 4.6	30 30 29 28 28 24 22 16 15 12	199 177 222 241 259 265 276 265 262 262	1. 0 1. 2 1. 4 2. 1 2. 6 3. 9 5. 9 5. 7 5. 3	31 31 26 24 22 19 17 16 15 15	292 312 290 291 237 230 239 249 255 254	2. 0 1. 3 1. 4 . 7 2. 2 2. 7 2. 9 4. 6 6. 1 8. 7	31 31 30 29 29 26 25 23 17 13	241 237 243 226 235 238 247 251 252 259 266 270	2.8 3.4 3.0 2.9 4.4 6.0 9.0 12.0 13.5 16.2 17.0 21.1	29 29 25 19 16 16 16 14 12 12	90 113 150 222 257 254 257 259 259 264 285	1. 2 2. 5 2. 3 2. 2 3. 1 4. 7 6. 2 5. 5 8. 4

MONTHLY WEATHER REVIEW

Table 3.—Maximum free air wind velocities (m. p. s.), for different sections of the United States, based on pilot-balloon observations during August 1940

		Surface	to 2,500	met	ers (m. s. l.)		Between 2,	500 and 8	5,000	meters (m. s. l.)		Abov	ve 5,000 ı	neter	s (m. s. l.)
Section	Maximum ve- locity	Direc- tion	Altitude (m.) m. s. l.	Date	Station	Maximum ve- locity	Direc- tion	Altitude (m.) m. s. l.	Date	Station	Maximum ve- locity	Direc- tion	Altitude (m.) m. s. l.	Date	Station
Northeast 1 East-Central 3 Southeast 3 North-Central 4 Central 5 South-Central 6 North-West 7 West-Central 8 Southwest 9 Southwest 9	35. 0 33. 0	WNW ESE E NNW SW SSW NW NNW	1,000 2,370 1,600 1,260 1,580 1,900 1,143 2,240 2,000	25 12 11 15 24 3 9 7	Caribou, Maine Knoxville, Tenn Spartansburg, S. C Minneapolis, Minn. Dodge City, Kans Amarillo, Tex Ellensburg, Wash Casper, Wyo El Paso, Tex	37. 1 28. S 21. 7 36. 6 33. 2 23. 2 37. 2 26. 5 26. 4	SSWESESWSWSWSWSW	3, 260 2, 501 2, 501 4, 600 3, 220 3, 910 3, 556 2, 501 2, 910	24 12 11 24 26 27 31 10 18	Elmira, N. Y. Knoxville, Tenn Spartansburg, S. C. Alpena, Mich Wichita, Kans. Oklahoma City, Okla. Havre, Mont. Cheyenne, Wyo El Paso, Tex.		WNW	16, 200 16, 840 33, 482 9, 540 12, 340 9, 510 7, 820 11, 395 13, 120	15 4 31 5 6 27 1 3 26	Nashville, Tenn. Miami, Fla. Huron, S. Dak.

Maine, Vermont, New Hampshire, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Pennsylvania and Northern Ohio.
 Delaware, Maryland, Virginia, West Virginia, Southern Ohio, Kentucky, Eastern Tennessee and North Carolina.
 South Carolina, Georgia, Florida and Alabama.
 Michigan, Wisconsin, Minnesota. North Dakota and South Dakota.
 Indiana, Illinois, Iowa, Nebraska, Kansas and Missouri.

Mississippi, Arkansas, Louisiana, Oklahoma, Texas (except extreme West Texas) and Western Tennessee.
 Montana, Idaho, Washington and Oregon.
 Wyoming, Colorado, Utah, Northern Nevada and Northern California.
 Southern California, Southern Nevada, Arizona, New Mexico, and extreme West Texas.

Table 4.—Mean altitudes and temperatures of significant points identifiable as tropopauses during August 1940, classified according to the potential temperatures (10° intervals between 290° and 409° A.) with which they are identified (based on radiosonde observations)

	Bisma	rck, N.	Dak.	Brow	nsville,	Tex.	Char	leston,	s. c.	Dei	ive r, C	olo.	E	ly, Nev		El	Paso, T	ex.	Great	Falls, I	Iont.
Potential tempera- tures, °A.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture °C.
290-299 300-309 310-319 320-329 330-339 340-349 350-359 360-369 370-379 380-389 300-399 400-409 Weighted means		9.0 10.7 11.9 13.0 13.8 14.7 15.0 15.4 16.9	-36. 1 -47. 6 -53. 1 -57. 3 -58. 8 -61. 2 -60. 1 -58. 0 -57. 8 -52. 7	1 12 14 13 10 8 4	11. 8 11. 8 13. 3 15. 0 16. 0 16. 4 16. 9 17. 2 14. 5	-56.0 -48.8 -57.7 -67.8 -72.7 -71.9 -72.0 -69.0 -63.3	2 14 18 10 6 5 3	11. 8 12. 3 13. 7 14. 8 15. 5 16. 3 16. 5	-57. 5 -54. 6 -62. 9 -67. 5 -69. 2 -71. 2 -69. 0	1 15 17 8 15 5 5 8 2 2	9.0 10.6 11.9 13.5 14.4 15.5 15.8 16.6 13.2	-36. 0 -44. 7 -51. 8 -61. 8 -63. 5 -68. 0 -65. 2 -65. 0 -56. 9	2 12 29 12 16 16 6 6	9. 7 10. 5 11. 8 13. 4 14. 4 15. 2 15. 0 16. 4 13. 3	-42.0 -44.3 -51.2 -60.2 -63.9 -65.6 -65.5 -63.8 -63.0 -57.4	3 18 11 13 9 2 4 5	9.8 12.1 13.8 14.7 15.7 16.4 16.8 17.1 14.1	-37. 3 -51. 8 -62. 5 -67. 3 -69. 8 -71. 0 -70. 8 -68. 2 -61. 6	11 32 14 28 4 5 4	9. 8 11. 2 12. 3 13. 8 14. 1 14. 3 15. 0 15. 8 16. 0 12. 3	-44. 1 -52. 3 -57. 6 -62. 5 -61. 5 -59. 5 -61. 2 -62. 7 -54. 9
Mean potential temperature °A. (weighted) No. days with ob- servations		352, 0 27			365, 0 23			359. 0 22			356. 6 23			358. 5 29			363.4 20			348. 7 30	
		Joliet, I	11.	Lak	ehurst,	N. J.	Me	dford, (Oreg.	Nas	hville, '	Tenn.	Oa	kland, (Calif.	Oklah	oma Ci	ty,Okla	. Оп	aha, N	br.
Potential temperatures °A.	Number of cases	Mean altitude (km.) m. s. l.	Mean tempera- ture °C.	Number of cases	Mean altitude (km.) m. s. l.	Mean tempera- ture °C.	Number of cases	Mean altitude (km.) m. s. l.	Mean tempera-	Number of cases	Mean altitude (km.) m. s. l.	Mean tempera- ture °C.	Number of cases	Mean altitude (km.) m. s. l.	Mean tempera- ture °C.	Number of cases	Mean altitude (km.) m. s. l.	Mean tempera- ture °C.	Number of cases	Mean altitude (km.) m. s. l.	Mean tempera- ture °C.
290-299 300-309	.	10.6											-			-	-	-			
310-319 320-329 330-339 340-349 350-359 360-369 370-379 380-389 390-399 400-409 Weighted means	. 11	10. 6 9. 7 10. 7 11. 8 13. 3 14. 3 15. 1 15. 4 16. 5 16. 2 13. 1	-44.5 -47.5 -52.3 -59.7 -63.0 -64.2 -63.9 -65.0 -60.0	3 9 24 7 8 6 8 3 1	9. 2 10. 5 12. 4 13. 2 14. 3 14. 7 15. 6 16. 1 16. 5 13. 1	-40. 3 -46. 2 -57. 0 -59. 6 -63. 9 -65. 5 -65. 3 -66. 0 -58. 1	1 9 14 3 2 3 7 3 3 3	10. 7 10. 8 12. 3 12. 6 13. 7 14. 9 15. 6 15. 8 16. 5 13. 3	-59.0	1 5 8 17 12 8 5 7 3	8. 7 10. 7 12. 7 13. 6 14. 6 15. 6 16. 1 16. 3 17. 0 14. 3	-47.8 -58.4 -61.9 -66.9 -68.4 -70.4 -67.6 -67.7	3 18 19 8 11 11 4 8 5	9. 4 10. 4 11. 7 13. 3 14. 3 15. 3 15. 8 16. 1 16. 5 13. 1	-44. 3 -44. 8 -50. 2 -59. 2 -67. 3 -67. 8 -67. 0 -64. 2 -56. 7	10 21 11 17 8 7 4 1	16. 6	-55. 0 -60. 6 -66. 8 -70. 4 -68. 7 -68. 8 -70. 0	19 9 9 7 6 7 3	9. 4 10. 2 12. 0 13. 3 14. 2 14. 6 15. 3 16. 1 16. 3 13. 0	-43.6 -43.0 -52.2 -59.7 -61.7 -61.4 -61.2 -63.9 -62.7 -55.4
Mean potential temperature A. (weighted)		355.7 24			355.7 29			359.4 15			363.8 25			358.2			358.6 24			358.3 25	

Table 4.—Mean altitudes and temperatures of significant points identifiable as tropopauses during August 1940, classified according to the potential temperatures (10° intervals between 290° and 409° A) with which they are identified (based on radiosonde obesrvations)—Continued

	Pho	enix, A	riz.	Sau	lt Ste. I Mich.	Marie,	Atlan	itic Sta.	No. 2 2		Pho	enix, Ar	iz.	Saul	t Ste. M Mich.	Iarie,	Atlan	tic Sta.	No. 2 2
Potential temperatures °A.	Num- ber of cases	Mean alti- tude (km.) m.s.l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- tude (km.) m.s.l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- tude (km.) m.s.l.	Mean tem- pera- ture °C.	Potential tempera- tures °A.	Num- ber of cases	Mean alti- tude (km.) m.s.l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- tude (km.) m.s.l.	Mean tem- pera- ture °C.	Num- ber of cases	Mean alti- ture (km.) m.s.l.	Mean tem- pera- ture °C.
290-299					7.3	-37. 0				390-399 400-409	9	16. 7	69.9	3	16. 2	-63.0	3	15.9	66. 3
310-319 320-329				3 10	8. 7 10. 0	-44.0 -49.5	1 1	8. 7 9. 7	-45.0 -49.0	Weighted means		14.3	-63.5		12.4	-54.9		13. 1	—59. 4
330–339 340–349 350–359 360–369 370–379 380–389	5 14 15 20 9 6	11. 6 12. 1 13. 7 14. 8 15. 4 16. 4	-56. 0 -51. 4 -61. 6 -67. 6 -68. 2 -72. 2	10 13 10 12 5 6	10.9 12.1 13.0 14.1 14.0 15.0	-51. 0 -54. 8 -57. 2 -62. 6 -57. 7 -60. 3	7 26 11 7 5	11. 5 12. 5 13. 5 14. 5 15. 0 15. 6	-53.7 -57.5 -61.6 -65.4 -64.8 -61.0	.0 .7 .5 Mean potential temperature °A. (weighted)		362.9 28			350.5 26			351.4 25	

² In or near the 5° square, lat. 40°00'N. to 45°00'N., long. 40°00'W. to 45°00'W.

WEATHER ON THE NORTH ATLANTIC OCEAN By H. C. Hunter

Atmospheric pressure.—The pressure was above normal, on the average, over most parts of the North Atlantic area which are covered by available reports. The excess was noteworthy over the west-central portion, where the Nantucket station reported a positive departure of 5.1 millibars (0.15 inch). The north-central West Indies region and the northern part of the Gulf of Mexico had pressure somewhat below normal.

The extremes of pressure in the vessel reports now at hand were 1,032.5 and 993.2 millibars (30.49 and 29.33 inches, respectively). The high mark was recorded on the American liner Extavia, near 40° N., 27° W., during the forenoon of the 5th. The low mark was noted not quite 100 miles to east-southeastward of Charleston, S. C., very early on the 11th, by the American steamship Tydolgas, which was then in the north semicircle of the second of the August tropical disturbances.

In some land areas affected by the first or the second of these cyclones it is noteworthy that stations near the coast recorded much lower readings than that of the *Tydolgas*, the lowest of these shore readings being 974.7 millibars (28.78 inches) at Savannah on the 11th.

Table 1.—Averages, departures, and extremes of atmospheric pressure (sea level) at selected stations for the North Atlantic Ocean and its shores, August 1940

Station	Average pressure	Depar- ture from normal	Highest	Date	Lowest	Date
Lisbon, Portugal Horta, Azores Belle Isle, Newfoundland Halifax, Nova Scotia Nantucket Hatteras Turks Island Key West New Orleans	Millibars 1, 018. 5 1, 023. 8 1, 014. 0 1, 020. 7 1, 017. 3 1, 015. 6 1, 015. 2 1, 014. 6	Millibars +1.9 +0.9 +1.8 +5.1 +1.4 -1.7 0.0 -0.6	Millibars 1, 023 1, 030 1, 029 1, 032 1, 031 1, 024 1, 017 1, 019 1, 020	17 9 23 30 11 12 3 2 30 31	Millibars 1, 011 1, 015 999 1, 010 1, 004 1, 005 1, 010 1, 012 1, 006	20 25, 26 26 20, 21 20 19 6 20 6

For 24 days.
For 20 days.

For those portions of the North Atlantic remote from the tropics the lowest reading found was 997.3 millibars (29.45 inches), noted on the United States Army transport American Legion, near the coast of northwestern Norway on the morning of the 18th.

Cyclones and gales.—On the ocean areas north of the 34th parallel of latitude there was little storminess worth mentioning. One occurrence of force 10 (whole gale) is indicated in the table of storms, this coming during the night of the 26th–27th at the location of the United States Coast Guard cutter Champlain, about 39° N.,59° W. A Low of considerable strength for the time of year had moved east-northeastward off the coast of Labrador during the preceding day, and a narrow trough extended southward from the center to about the 35th parallel of latitude, causing strong shifting winds within a small strip.

Disturbances of the tropical region.—Three cyclones affected the North Atlantic waters within the tropics and a short distance outside during the month. The first two of these are described elsewhere in this Review, but the third storm is to be described in the next succeeding issue.

The earliest of these storms formed east of Florida on the 2d or 3d and moved slowly to the coast of southwestern Louisiana by the 7th, seemingly never reaching hurricane intensity, though one vessel met a force 11 wind.

The second storm traveled northwestward from the vicinity of St. Thomas on the 5th to the coast of South Carolina on the 11th. During the final hours of its passage over water there were hurricane force winds, as noted by two vessels. The tracks of the first and second storms are indicated on an accompanying map.

The third storm, coming from the southeastward, was not far off Hatteras when the month ended.

Fog.—Reports indicate much less fog than normal, especially near New England and Nova Scotia. However, the foggiest 5° square for the entire North Atlantic ocean area was in this section, 40° to 45° N., 65° to 70° W., where fog was observed on 5 days, the normal August occurrence there being 17 days.

During the 9-day period, 9th to 17th, there was no fog in any part of the ocean that available reports were covering on those days.

² Also 5 later dates.

Note.—All data based on available observations, departures compiled from best available normals related to time of observation, except Hatteras, Key West, Nantucket, and New Orleans, which are 24-hour corrected means.